



Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

J. Math. Anal. Appl. 297 (2004) 267–284

Journal of
MATHEMATICAL
ANALYSIS AND
APPLICATIONS

www.elsevier.com/locate/jmaa

Periodic and subharmonic solutions of a class of superquadratic Hamiltonian systems [☆]

Shang-Jie Chen, Chun-Lei Tang ^{*}

Department of Mathematics, Southwest Normal University, Chongqing 400715, People's Republic of China

Received 12 February 2004

Available online 8 July 2004

Submitted by H.L. Smith

Abstract

Periodic solutions and infinitely distinct subharmonic solutions are obtained for a class of nonconvex and nonautonomous superquadratic Hamiltonian systems $\dot{z} = \mathcal{J}H_z(z, t)$ by using the minimax methods in critical point theory.

© 2004 Elsevier Inc. All rights reserved.

Keywords: Periodic solutions; Subharmonic solutions; Superquadratic Hamiltonian systems; Minimax method; Truncating argument; Critical point

1. Introduction and main results

Consider the nonautonomous Hamiltonian system

$$\dot{z} = \mathcal{J}H_z(z, t), \quad (1.1)$$

where

$$\mathcal{J} = \begin{pmatrix} 0 & -I_n \\ I_n & 0 \end{pmatrix}$$

[☆] Supported by National Natural Science Foundation of China, by Major Project of Science and Technology of MOE, PR China, and by the Teaching and Research Award Program for Outstanding Young Teachers in Higher Education Institutions of MOE, PR China.

^{*} Corresponding author.

E-mail address: tangcl@swnu.edu.cn (C.-L. Tang).

is the standard symplectic matrix, $H : \mathcal{R}^{2n} \times \mathcal{R} \rightarrow \mathcal{R}$ is a C^1 function, and T -periodic in the second variable. A usual hypothesis under which interesting results for (1.1) can be obtained is

(C) There exist $\mu > 2$ and $r_0 > 0$ such that

$$\frac{1}{\mu} H_z(z, t) \cdot z \geq H(z, t) > 0 \quad \forall z \in \mathcal{R}^{2n}, |z| \geq r_0, \forall t \in \mathcal{R}.$$

Here, and in what follows, we denote by \cdot the usual inner product in \mathcal{C}^m and by $|\cdot|$ the corresponding norm. A Hamiltonian satisfying (C) is called superquadratic. There are several papers dealing with the existence of T -periodic solutions of (1.1) under condition (C) (see, e.g., [1–10, 17]).

A particular case of Eq. (1.1) is the so-called second order Hamiltonian system

$$\ddot{q} + \nabla V(q, t) = 0, \quad (1.2)$$

where $V : \mathcal{R}^n \times \mathcal{R} \rightarrow \mathcal{R}$ is a C^1 function, and T -periodic in the second variable. If we put

$$H(p, q, t) = \frac{1}{2}|p|^2 + V(q, t), \quad (1.3)$$

(1.2) is equivalent to (1.1). Several results for the existence of T -periodic solutions for (1.2) have been obtained. A usual hypothesis to consider on V is

(2C) There exist $\mu > 2$ and $r_0 > 0$ such that

$$\frac{1}{\mu} \nabla V(q, t) \cdot q \geq V(q, t) > 0 \quad \forall q \in \mathcal{R}^n, |q| \geq r_0, \forall t \in \mathcal{R}.$$

See, for example, [8, 11–13] and others. We observe that H does not satisfy (C) if V satisfies (2C) and H is defined by (1.3).

In 1993, Felmer [1] extend some existence result for (1.1) where the Hamiltonian satisfies a superquadratic condition that include simultaneously (C) and (2C). In [1], Felmer gave the following condition:

(3C) There exist $\alpha > 1$, $\beta > 1$, $1/\alpha + 1/\beta < 1$ and $r_0 > 0$ such that

$$\begin{aligned} \frac{1}{\alpha} H_p(p, q, t) \cdot p + \frac{1}{\beta} H_q(p, q, t) \cdot q &\geq H(p, q, t) > 0 \\ \forall z = (p, q) \in \mathcal{R}^n \times \mathcal{R}^n, |z| &\geq r_0, \forall t \in \mathcal{R}. \end{aligned}$$

H obviously satisfies (3C) when it satisfies (C). It is easy to obtain that when V satisfies (2C) and H is defined by (1.3), H satisfies (3C) too.

Felmer [1] had well solved the unifying of superquadratic conditions of Eqs. (1.1) and (1.2), but the result of Felmer in [1] does not include the corresponding result in [7] as a special case. A natural question is whether there exists a result which contains the corresponding results in [1] and [7] as a special case.

Motivated by [1] and [7], we give this question a positive answer by the minimax methods in the critical point theory and obtain a result (see Theorem 1.1) unifies and generalizes Theorem 0.1 in [1] and Theorem 2.49 in [7]. In this paper, we consider a Hamiltonian satisfying the following hypothesis:

(H0) H is of class C^1 , and

$$H(z, t + T) = H(z, t) \quad \forall z \in \mathbb{R}^{2n}, \quad \forall t \in \mathbb{R},$$

(H1) $H(z, t) \geq 0 \quad \forall t \in \mathbb{R}, \quad \forall z \in \mathbb{R}^{2n}$,

(H2) $\exists \alpha > 1, \beta > 1, 1/\alpha + 1/\beta < 1$ and $r_0 > 0$ such that

$$\frac{1}{\alpha} H_p(p, q, t) \cdot p + \frac{1}{\beta} H_q(p, q, t) \cdot q \geq H(p, q, t) > 0$$

$$\forall z = (p, q) \in \mathbb{R}^{2n}, \quad |z| \geq r_0, \quad \forall t \in \mathbb{R},$$

(H3) $\lim_{|(p,q)| \rightarrow 0} \frac{H(p, q, t)}{|p|^{1+\alpha/\beta} + |q|^{1+\beta/\alpha}} = 0$,

(H4) $\exists a > 0$ and $b \geq 0$ such that

$$|H_z(p, q, t)| \leq a \left(\frac{1}{\alpha} H_p(p, q, t) \cdot p + \frac{1}{\beta} H_q(p, q, t) \cdot q \right) + b$$

$$\forall z = (p, q) \in \mathbb{R}^{2n}, \quad \forall t \in \mathbb{R}.$$

We obtain the following theorem.

Theorem 1.1. *Suppose that H satisfies (H0)–(H4). Then Eq. (1.1) has a nonzero T -periodic solution.*

Remark 1.1. It is easy to obtain that our Theorem 1.1 unifies and generalizes Theorem 0.1 in [1], Theorem 2.49 in [7] and Corollary 1.5 in [5]. Even if it is in the autonomous case, there are functions H satisfying our Theorem 1.1 and not satisfying the corresponding results in [1,7] and others. In fact, for $\alpha > 1, \beta > 1$ satisfying $1/\alpha + 1/\beta < 1$, let

$$H(z) = a_1(|p|^{1+\alpha/\beta} + |q|^{1+\beta/\alpha})^{\gamma_1} + a_2(|p|^{1+\alpha/\beta} + |q|^{1+\beta/\alpha})^{\gamma_2}, \quad (1.4)$$

where $a_1 > 0, a_2 > 0, 1 < \gamma_1 < \alpha\beta/(\alpha + \beta) < \gamma_2$. Then H satisfies the conditions of our Theorem 1.1, but does not satisfy the corresponding theorems in [1–10,17].

In the following, we search for kT periodic solutions (called subharmonics) of (1.1) and (1.2). Several results for the existence of subharmonic solutions for (1.1) have been obtained when H is convex (see [5,14–16,18,19]). There are also several papers dealing with the existence of subharmonic solutions of (1.1) when H is nonconvex (see [5,10,20–22]). In [5,10,21–23], subharmonic solutions are obtained for superquadratic Hamiltonian systems. In [5,20], subharmonic solutions are obtained for subquadratic Hamiltonian systems. Many results for the existence of subharmonic solutions for (1.2) have been obtained (see [5,14,18,24–31]). Here, we give the following theorem.

Theorem 1.2. *Suppose that H satisfies (H0)–(H4). Then (1.1) possesses infinitely distinct subharmonic solutions.*

Remark 1.2. Theorem 1.2 generalizes Theorem 1.36 of [5] in the case that the quadratic form vanishes. It is easy to obtain that the functions H in (1.4) satisfy the conditions of our Theorem 1.2, but do not satisfy that of the corresponding theorems in [5, 14–16, 18–23].

2. Proofs of the main results

We study the existence of T -periodic solutions of (1.1) from the variational point of view. Taking $\omega = 2\pi/T$, let $E := W^{1/2,2}([0, T], \mathcal{R}^{2n})$ be the Sobolev's space of T -periodic \mathcal{R}^{2n} -valued functions

$$z(t) = \sum_{j \in \mathbb{Z}} a_j e^{\omega i j t}, \quad a_{-j} = \bar{a}_j \in \mathbb{C}^{2n},$$

such that

$$\|z\|^2 = \frac{T}{2} \sum_{j \in \mathbb{Z} - \{0\}} |j| |a_j|^2 + T |a_0|^2 < \infty.$$

An inner product in E is defined by

$$\langle z, \eta \rangle = \frac{T}{2} \sum_{j \in \mathbb{Z} - \{0\}} |j| a_j \cdot \bar{b}_j + T a_0 \cdot \bar{b}_0.$$

It is well known that E compactly embedded in $L^\gamma([0, T], \mathcal{R}^{2n}) \equiv L^\gamma$ for $\gamma \in [1, +\infty)$ and as a consequence there exists a constant $C > 0$ such that

$$\|z\|_\gamma \leq C \|z\| \quad \forall z \in E \quad (2.1)$$

for

$$\gamma = 1, \alpha, \beta, 1 + \frac{\alpha}{\beta}, 1 + \frac{\beta}{\alpha}.$$

Here and in what follows $\|\cdot\|_\gamma$ denotes the usual norm in L^γ . For $z = (p, q)$ and $\eta = (\phi, \psi)$ in E and smooth we define

$$B(z, \eta) = \int_0^T (p \cdot \dot{\psi} + \phi \cdot \dot{q}) dt, \quad A(z) = \frac{1}{2} B(z, z). \quad (2.2)$$

Both A and B can be extended continuously to the whole space E , and the bilinear form B induces a linear, bounded, selfadjoint operator $L : E \rightarrow E$ defined by

$$B(z, \eta) = \langle Lz, \eta \rangle \quad \forall z, \eta \in E. \quad (2.3)$$

In E we can consider the splitting

$$E = E_+ \oplus E_- \oplus E_0, \quad (2.4)$$

where

$$\begin{aligned} E_+ &= \overline{\text{span}}^E \left\{ \sin(\omega j t) e_k - \cos(\omega j t) e_{k+n}, \right. \\ &\quad \left. \cos(\omega j t) e_k + \sin(\omega j t) e_{k+n} \mid j \in \mathcal{N}, 1 \leq k \leq n \right\}, \\ E_- &= \overline{\text{span}}^E \left\{ \sin(\omega j t) e_k + \cos(\omega j t) e_{k+n}, \right. \\ &\quad \left. \cos(\omega j t) e_k - \sin(\omega j t) e_{k+n} \mid j \in \mathcal{N}, 1 \leq k \leq n \right\}, \end{aligned}$$

and

$$E_0 = \text{span}\{e_1, \dots, e_{2n}\}.$$

Here $\{e_1, \dots, e_{2n}\}$ is the canonical basis for \mathcal{R}^{2n} . We observe that A is positive on E_+ , negative on E_- , and it vanishes on E_0 . Actually E_+ , E_- , and E_0 are the positive, negative, and null eigenspace of the linear operator L , respectively. On the space E we can see that

$$\|z\|^2 = A(z_+) - A(z_-) + T|z_0|^2, \quad (2.5)$$

where $z = z_+ + z_- + z_0$ with $z_+ \in E_+$, $z_- \in E_-$ and $z_0 \in E_0$.

Formally the T -periodic solutions of (1.1) are the critical points of the functional

$$J(z) = A(z) - \int_0^T H(z, t) dt.$$

However since we do not have an adequate control on the growth of H , J is not needed to be well defined. To overcome this difficulty we use a truncating argument introduced by Rabinowitz [7]. Let $K \geq 1$ be a constant and $\chi_K \in C^\infty(\mathcal{R}, \mathcal{R})$ such that $\chi_K(s) \equiv 1$ if $s \leq K$, $\chi_K(s) \equiv 0$ if $s \geq K+1$, and $\chi'_K(s) < 0$ for $s \in (K, K+1)$. We define

$$H_K(z, t) = \chi_K(|z|)H(z, t) + (1 - \chi_K(|z|))r_K(|p|^\alpha + |q|^\beta),$$

where

$$r_K = \max \left\{ \frac{H(z, t)}{|p|^\alpha + |q|^\beta} : K \leq |z| \leq K+1, t \in \mathcal{R} \right\}.$$

An easy computation shows that H_K satisfies the same hypothesis as H . Moreover, hypothesis (H2) and (H4) are satisfied with constants independent of K . With this change the functional

$$J_K(z) = A(z) - \int_0^T H_K(z, t) dt$$

is well defined on E and it is of class C^1 . Next we show a lemma regarding the growth of H as a consequence of hypothesis (H0) and (H2) and hence also valid for H_K .

Lemma 2.1 [1]. *If H satisfies (H0) and (H2), there exist constants $c_1 \geq 0$ and $c_2 \geq 0$ such that*

$$H(p, q, t) \geq c_1(|p|^\alpha + |q|^\beta) - c_2 \quad \forall (p, q) \in \mathcal{R}^{2n}, \forall t \in \mathcal{R}.$$

For needs of our proofs, we introduce the following abstract theorem due to Felmer [1].

We consider a Hilbert space E with inner product $\langle \cdot, \cdot \rangle$ and norm $\| \cdot \|$. We assume that E has a splitting $E = X \oplus Y$, where the subspace X and Y are not necessarily orthogonal and both of them can be infinite dimensional. Let $I : E \rightarrow \mathbb{R}$ be a functional having the structure

$$I(z) = \langle Lz, z \rangle + b(z).$$

(I1) $L : E \rightarrow E$ is a linear, bounded, selfadjoint operator,

(I2) b' is compact,

there are two linear bounded, invertible operators $B_1, B_2 : E \rightarrow E$ satisfying

(I3) if $v \in \mathcal{R}_+$, the linear operator

$$\hat{B}(v) = P_X B_1^{-1} \exp(vL) B_2 : X \rightarrow X$$

is invertible. Here P_X denotes the projection of E onto X inducing by the splitting $E = X \oplus Y$, and \mathcal{R}_+ is a set of nonnegative real numbers.

Let $\rho > 0$ and define

$$S = \{B_1 z \mid \|z\| = \rho, z \in Y\}. \quad (2.6)$$

For $z_+ \in Y$, $z_+ \neq 0$, $\sigma > \rho / \|B_1^{-1} B_2 z_+\|$, and $M > 0$, we define

$$Q = \{B_2(s z_+ + z) \mid 0 \leq s \leq \sigma, \|z\| \leq M, z \in X\}. \quad (2.7)$$

We define ∂Q as the boundary of Q relative to the subspace $\{B_2(s z_+ + z) \mid s \in \mathcal{R}, z \in X\}$. Let us consider the class of functions

$$\Gamma = \{h \in C(E \times [0, 1], E) \mid h \text{ satisfies } \Gamma_1, \Gamma_2, \text{ and } \Gamma_3\},$$

where

(Γ_1) $h(z, t) = \exp(v(z, t)L)z + \tilde{K}(z, t)$, where $v : E \times [0, 1] \rightarrow \mathcal{R}^+$ is continuous and transforms bounded sets into bounded sets, and $\tilde{K} : E \times [0, 1] \rightarrow E$ is compact,

(Γ_2) $h(z, t) = z \forall z \in \partial Q$,

(Γ_3) $h(z, 0) = z \forall z \in Q$.

Theorem 2.2 [1]. *Let $I : E \rightarrow \mathbb{R}$ be a C^1 functional satisfying the Palais–Smale condition and (I1)–(I3). Furthermore assume that there is a constant $\delta > 0$ such that*

(i) $I(z) \geq \delta \forall z \in S$,

(ii) $I(z) \leq 0 \forall z \in \partial Q$.

Then I possesses a critical point with critical value $d \geq \delta$ characterized by

$$d = \inf_{h \in \Gamma} \sup_{z \in Q} I(h(z, 1)).$$

Before proving our theorems, we obtained some useful facts regarding the subspaces E_+ and E_- and some estimates. Given a function $z \in L^\gamma$, $\gamma \geq 1$, with Fourier series given by

$$z = \sum_{k \geq 1} a_k \cos \omega k t + b_k \sin \omega k t,$$

with $a_k, b_k \in \mathcal{R}^n$, we define the conjugate of z by

$$\bar{z} = \sum_{k \geq 1} b_k \cos \omega k t - a_k \sin \omega k t.$$

We note that $\bar{\bar{z}} = -z \ \forall z \in L^\gamma$.

Lemma 2.3 [1]. *An element $z = (p, q) \in E$ belongs to E_+ (respectively E_-) if and only if $\bar{p} = -q$ (respectively $\bar{p} = q$).*

Lemma 2.4 [1]. *For every $z = (p, q) \in E_+$ or E_- , it holds that*

$$\|p\| = \|q\| = \frac{1}{\sqrt{2}} \|z\|.$$

Lemma 2.5 [1]. *Let $P_+ : E \rightarrow E$ (respectively $P_- : E \rightarrow E$) be projections induced by the splitting $E = E_+ \oplus E_- \oplus E_0$. If $z = (p, q) + z_0$, $z_0 \in E_0$, one has*

$$P_+(z) = \frac{1}{2}(p + \bar{q}, q - \bar{p}) \quad \left(\text{respectively } P_-(z) = \frac{1}{2}(p - \bar{q}, q + \bar{p}) \right).$$

We define the operator B_1 and B_2 and the splitting E . Let $X = E_- + E_0$ and $Y = E_+$. We define $B_1 : E \rightarrow E$ by

$$B_1(z) = B_1((p, q)) = (\rho^{\beta-1} p, \rho^{\alpha-1} q), \quad (2.8)$$

and $B_2 : E \rightarrow E$ by

$$B_2(z + z_0) = B_2((p, q) + z_0) = (\sigma^{\beta-1} p, \sigma^{\alpha-1} q) + z_0, \quad (2.9)$$

where $z_0 \in E_0$ and $z = (p, q) \in E_- + E_+$, and the constants ρ and δ will be defined in Lemmas 2.7 and 2.8. Certainly B_1 and B_2 are bounded linear operators and both of them are invertible. From (2.6) and (2.8), we obtain

$$S = \{(\rho^{\beta-1} p, \rho^{\alpha-1} q) \mid \|(p, q)\| = \rho, (p, q) \in E_+\}. \quad (2.10)$$

By (2.7) and (2.9), we have

$$Q = \{s(\sigma^{\beta-1} p_+, \sigma^{\alpha-1} q_+) + (\sigma^{\beta-1} p, \sigma^{\alpha-1} q) + z_0 \mid 0 \leq s \leq \sigma, 0 \leq \|(p, q) + z_0\| \leq M, (p, q) \in E_-, z_0 \in E_0\}, \quad (2.11)$$

where $z_+ = (p_+, q_+) \in E_+$ is a fixed eigenvector of L associated to an eigenvalue $\lambda > 0$ and $\|z_+\| = 1$. In what follows we denote by ∂Q the boundary of Q relative to the subspace

$$\{s(\sigma^{\beta-1} p_+, \sigma^{\alpha-1} q_+) + (\sigma^{\beta-1} p, \sigma^{\alpha-1} q) + z_0 \mid s \in \mathcal{R}, (p, q) \in E_-, z_0 \in E_0\}.$$

Lemma 2.6 [1]. *The functional J_K satisfies the Palais–Smale condition.*

Lemma 2.7. *There exist $\rho > 0$ such that (i) is satisfied for J_K when S is defined by (2.10).*

Proof. From hypothesis (H3) and the form of H_K we have for each $\varepsilon > 0$,

$$H_K(p, q, t) \leq \varepsilon(|p|^{1+\alpha/\beta} + |q|^{1+\beta/\alpha}) + c_3(|p|^\alpha + |q|^\beta), \quad (2.12)$$

where $c_3 = c_3(\varepsilon, K) > 0$. Let $(p, q) \in E_+$ and take $z = (\rho^{\beta-1}p, \rho^{\alpha-1}q)$ for some $\rho > 0$. Then from (2.12), we have

$$\begin{aligned} J_K(z) &\geq \rho^{\alpha+\beta-2} \|(p, q)\|^2 - \varepsilon(\rho^{(\beta-1)(1+\alpha/\beta)} \|p\|_{1+\alpha/\beta}^{1+\alpha/\beta} + \rho^{(\alpha-1)(1+\beta/\alpha)} \|q\|_{1+\beta/\alpha}^{1+\beta/\alpha}) \\ &\quad - c_3(\rho^{(\beta-1)\alpha} \|p\|_\alpha^\alpha + \rho^{(\alpha-1)\beta} \|q\|_\beta^\beta). \end{aligned} \quad (2.13)$$

From (2.13) and (2.1) we obtain

$$\begin{aligned} J_K(z) &\geq \rho^{\alpha+\beta-2} \|(p, q)\|^2 - \varepsilon c_4(\rho^{(\beta-1)(1+\alpha/\beta)} \|p\|^{1+\alpha/\beta} \\ &\quad + \rho^{(\alpha-1)(1+\beta/\alpha)} \|q\|^{1+\beta/\alpha}) - c_5(\rho^{(\beta-1)\alpha} \|p\|^\alpha + \rho^{(\alpha-1)\beta} \|q\|^\beta), \end{aligned} \quad (2.14)$$

where $c_4 = \max\{C^{1+\alpha/\beta}, C^{1+\beta/\alpha}\}$, $c_5 = c_3 \max\{C^\alpha, C^\beta\}$. If we consider $\|(p, q)\| = \rho$, from Lemma 2.4 and (2.14) one obtains

$$\begin{aligned} J_K(z) &\geq \rho^{\alpha+\beta} - \varepsilon c_4 \rho^{\alpha+\beta} \left(\left(\frac{1}{\sqrt{2}} \right)^{1+\alpha/\beta} + \left(\frac{1}{\sqrt{2}} \right)^{1+\beta/\alpha} \right) \\ &\quad - c_5 \rho^{\alpha\beta} \left(\left(\frac{1}{\sqrt{2}} \right)^\alpha + \left(\frac{1}{\sqrt{2}} \right)^\beta \right). \end{aligned} \quad (2.15)$$

Noticing $1/\sqrt{2} < 1$, then we have from (2.15),

$$J_K(z) \geq (1 - 2\varepsilon c_4) \rho^{\alpha+\beta} - 2c_5 \rho^{\alpha\beta}.$$

Taking $\varepsilon = 1/(4c_4)$, and then, since α and β satisfy $1/\alpha + 1/\beta < 1$, there exist $\rho > 0$ and $\delta > 0$ such that

$$J_K(z) \geq \delta > 0$$

and this inequality holds for $z \in S$, according to the definition of S . \square

Lemma 2.8. *There are constants $\sigma > 0$ and $M > 0$ such that J_K satisfies (ii) for Q defined by (2.11).*

Proof. For $s \in \mathcal{R}^+$, $(p, q) \in E_-$, and $z_0 = (p_0, q_0) \in E_0$ we take

$$z = s(\sigma^{\beta-1}p_+, \sigma^{\alpha-1}q_+) + (\sigma^{\beta-1}p, \sigma^{\alpha-1}q) + z_0.$$

Then we have

$$\begin{aligned}
A(z) &= \frac{1}{2} B(z, z) = \int_0^T (s\sigma^{\beta-1} p_+ + \sigma^{\beta-1} p + p_0) \cdot (s\sigma^{\alpha-1} \dot{q}_+ + \sigma^{\alpha-1} \dot{q}) dt \\
&= \int_0^T \sigma^{\alpha+\beta-2} (s^2 p_+ \cdot \dot{q}_+ + p \cdot \dot{q}) dt \\
&= \sigma^{\alpha+\beta-2} \left(\frac{1}{2} s^2 B(z_+, z_+) + A((p, q)) \right) \\
&= \sigma^{\alpha+\beta-2} \left(\frac{1}{2} s^2 \langle Lz_+, z_+ \rangle - \|(p, q)\|^2 \right) \\
&= \sigma^{\alpha+\beta-2} \left(\frac{1}{2} s^2 \langle \lambda z_+, z_+ \rangle - \|(p, q)\|^2 \right) \\
&= \sigma^{\alpha+\beta-2} \left(\frac{1}{2} s^2 \lambda - \|(p, q)\|^2 \right)
\end{aligned} \tag{2.16}$$

by (2.2), (2.4), (2.3), (2.5) and $z_+ = (p_+, q_+) \in E_+$ is a fixed eigenvector of L associated to an eigenvalue λ and $\|z_+\| = 1$. From hypothesis (H1) we see that for $s = 0$,

$$J_K(z) \leq 0. \tag{2.17}$$

Since $\int_0^T \sin \omega j t dt = 0$, $\int_0^T \cos \omega j t dt = 0$, $(p_+, q_+) \in E_+$, $(p, q) \in E_-$, and definition of E_+ , E_- we obtain

$$\int_0^T (\sigma^{\beta-1}(sp_+ + p) + p_0) dt = \int_0^T p_0 dt = T \cdot p_0. \tag{2.18}$$

It follows from (2.18) and Hölder inequality that

$$\begin{aligned}
|p_0| &= \frac{1}{T} \left| \int_0^T (\sigma^{\beta-1}(sp_+ + p) + p_0) dt \right| \\
&\leq \frac{1}{T} \int_0^T |\sigma^{\beta-1}(sp_+ + p) + p_0| dt \\
&\leq \frac{1}{T} \left(\int_0^T |\sigma^{\beta-1}(sp_+ + p) + p_0|^\alpha dt \right)^{1/\alpha} \cdot \left(\int_0^T 1^{\alpha/(\alpha-1)} dt \right)^{(\alpha-1)/\alpha} \\
&= (T)^{-1/\alpha} \|\sigma^{\beta-1}(sp_+ + p) + p_0\|_\alpha.
\end{aligned}$$

Then, we have

$$T |p_0|^\alpha \leq \|\sigma^{\beta-1}(sp_+ + p) + p_0\|_\alpha^\alpha. \tag{2.19}$$

From (2.19) we obtain

$$\begin{aligned}
\sigma^{(\beta-1)\alpha} \| (sp_+ + p) \|_\alpha^\alpha &= \int_0^T |\sigma^{\beta-1}(sp_+ + p)|^\alpha dt \\
&\leq 2^\alpha \int_0^T (|\sigma^{\beta-1}(sp_+ + p) + p_0|^\alpha + |p_0|^\alpha) dt \\
&= 2^\alpha (\| \sigma^{\beta-1}(sp_+ + p) + p_0 \|_\alpha^\alpha + T |p_0|^\alpha) \\
&\leq 2^{\alpha+1} \| \sigma^{\beta-1}(sp_+ + p) + p_0 \|_\alpha^\alpha.
\end{aligned} \tag{2.20}$$

By (2.19) and (2.20), we have

$$\begin{aligned}
&\sigma^{(\beta-1)\alpha} \| (sp_+ + p) \|_\alpha^\alpha + T |p_0|^\alpha \\
&\leq 2^{\alpha+1} \| \sigma^{\beta-1}(sp_+ + p) + p_0 \|_\alpha^\alpha + \| \sigma^{\beta-1}(sp_+ + p) + p_0 \|_\alpha^\alpha \\
&\leq 2^{\alpha+2} \| \sigma^{\beta-1}(sp_+ + p) + p_0 \|_\alpha^\alpha,
\end{aligned}$$

i.e.,

$$\| \sigma^{\beta-1}(sp_+ + p) + p_0 \|_\alpha^\alpha \geq \frac{1}{2^{\alpha+2}} (\sigma^{(\beta-1)\alpha} \| (sp_+ + p) \|_\alpha^\alpha + T |p_0|^\alpha). \tag{2.21}$$

Similarly, one has

$$\| \sigma^{\alpha-1}(sq_+ + q) + q_0 \|_\beta^\beta \geq \frac{1}{2^{\beta+2}} (\sigma^{(\alpha-1)\beta} \| (sq_+ + q) \|_\beta^\beta + T |q_0|^\beta). \tag{2.22}$$

Taking $c_6 = c_1 \min\{1/2^{\alpha+2}, 1/2^{\beta+2}\}$, for every point in Q we have from Lemma 2.1, (2.21) and (2.22),

$$\begin{aligned}
\int_0^T H_K(z, t) dt &\geq c_1 (\| \sigma^{\beta-1}(sp_+ + p) + p_0 \|_\alpha^\alpha + \| \sigma^{\alpha-1}(sq_+ + q) + q_0 \|_\beta^\beta) - T c_2 \\
&\geq c_6 (\sigma^{(\beta-1)\alpha} \| sp_+ + p \|_\alpha^\alpha + \sigma^{(\alpha-1)\beta} \| sq_+ + q \|_\beta^\beta \\
&\quad + T (|p_0|^\alpha + |q_0|^\beta)) - T c_2.
\end{aligned} \tag{2.23}$$

We assume now that $\alpha \leq \beta$ (the case $\alpha \geq \beta$ can be treated in an analogous way). By the triangle inequality we have

$$\| sp_+ \|_\alpha \leq \frac{1}{2} \| sp_+ + p \|_\alpha + \frac{1}{2} \| sp_+ - p \|_\alpha. \tag{2.24}$$

From Lemma 2.3 and the fact that for all $z \in L^\gamma$, $\bar{\bar{z}} = -z$, one has for $(p, q) \in E_-$,

$$\bar{p} = q, \quad \bar{\bar{p}} = \bar{q}, \quad -p = \bar{q}. \tag{2.25}$$

Similarly, for $(p_+, q_+) \in E_+$ we obtain

$$p_+ = \bar{q}_+. \tag{2.26}$$

From (2.25), (2.26) and Marchel Riesz's theorem in Section 12.9 of [32], one obtains

$$\| sp_+ - p \|_\alpha = \| s\bar{q}_+ + \bar{q} \|_\alpha \leq C_0 \| sq_+ + q \|_\alpha. \tag{2.27}$$

Using Hölder inequality, we have

$$\begin{aligned} \|sq_+ + q\|_\alpha^\alpha &= \int_0^T |sq_+ + q|^\alpha \cdot 1 \, dt \\ &\leq \left(\int_0^T |sq_+ + q|^{\alpha \cdot (\beta/\alpha)} \, dt \right)^{\alpha/\beta} \cdot \left(\int_0^T 1^{\beta/(\beta-\alpha)} \, dt \right)^{(\beta-\alpha)/\beta} \\ &= \|sq_+ + q\|_\beta^\alpha \cdot T^{(\beta-\alpha)/\beta}. \end{aligned} \quad (2.28)$$

Taking $c_7 = \frac{1}{2}C_0T^{(\beta-\alpha)/(\alpha\beta)}$, from (2.24), (2.27) and (2.28) we obtain

$$\|sp_+ \|_\alpha \leq \frac{1}{2}\|sp_+ + p\|_\alpha + c_7\|sq_+ + q\|_\beta, \quad (2.29)$$

and then the existence of a constant $c > 0$ for all $s > 0$, $(p, q) \in E_-$ follows such that

$$\|sp_+ + p\|_\alpha \geq sc \quad (2.30)$$

or

$$\|sq_+ + q\|_\beta \geq sc. \quad (2.31)$$

Otherwise, for every $c > 0$, there exists $s > 0$ and $(p, q) \in E_-$ such that

$$\|sp_+ + p\|_\alpha \leq sc, \quad \|sq_+ + q\|_\beta \leq sc. \quad (2.32)$$

Then, from (2.29) and (2.32) we have

$$\|sp_+ \|_\alpha \leq sc \left(\frac{1}{2} + c_4 \right).$$

Thus, $\|p_+ \|_\alpha \leq (1/2 + c_4)c$. Since c is random, we obtain $p_+ = 0$. And we have $q_+ = 0$ from Lemma 2.4. It contradicts that $\|(p_+, q_+)\| = 1$. In case that (2.30) holds, it follows from (2.16) and (2.23) that

$$J_K(z) \leq \frac{1}{2}\sigma^{\alpha+\beta-2}\lambda s^2 - c_6c^\alpha\sigma^{(\beta-1)\alpha}s^\alpha + Tc_2. \quad (2.33)$$

In case (2.31) holds, it follows from (2.16) and (2.23) that

$$J_K(z) \leq \frac{1}{2}\sigma^{\alpha+\beta-2}\lambda s^2 - c_6c^\beta\sigma^{(\alpha-1)\beta}s^\beta + Tc_2. \quad (2.34)$$

Choosing $s = \sigma$, and taking σ large enough it follows from $1/\alpha + 1/\beta < 1$, (2.33) and (2.34) that

$$J_K(z) \leq 0. \quad (2.35)$$

Finally we choose M . Given $s \in (0, \sigma)$ we have from (2.16) and (2.23) that

$$\begin{aligned} J_K(z) &\leq \frac{1}{2}\sigma^{\alpha+\beta-2}\lambda s^2 - \sigma^{\alpha+\beta-2}\|(p, q)\|^2 - Tc_6(|p_0|^\alpha + |q_0|^\beta) - Tc_2 \\ &\leq \frac{1}{2}\lambda\sigma^{\alpha+\beta} - \sigma^{\alpha+\beta-2}\|(p, q)\|^2 - Tc_6(|p_0|^\alpha + |q_0|^\beta) - Tc_2, \end{aligned}$$

so that if M is enough large and $\|(p, q) + z_0\| = M$ we have

$$J_K(z) \leq 0. \quad (2.36)$$

Thus, from (2.17), (2.35) and (2.36) we obtain

$$J_K(z) \leq 0 \quad \forall z \in \partial Q. \quad \square$$

Lemma 2.9. J_K satisfies (I1)–(I3).

Proof. From (2.1) and (2.2) we have

$$\begin{aligned} J_K(z) &= A(z) - \int_0^T H_K(z, t) dt = \frac{1}{2} B(z, z) - \int_0^T H_K(z, t) dt \\ &= \frac{1}{2} \langle Lz, z \rangle - \int_0^T H_K(z, t) dt. \end{aligned}$$

Taking $\Psi_K(z) = \int_0^T H_K(z, t) dt$, then for all $u \in E$ we obtain

$$(J'_K(z), u) = \langle Lz, u \rangle - \int_0^T \nabla H_K(z, t) \cdot u dt = \langle Lz, u \rangle - \langle \nabla \Psi_K(z), u \rangle.$$

So $\nabla J_K = L + (-\nabla \Psi_K)$, where L is a linear bounded selfadjoint operator and $(-\nabla \Psi_K)$ is a compact operator. Thus, J_K satisfies (I1) and (I2).

From the definition of L , one has

$$\langle Lz, \eta \rangle = \int_0^T p \cdot \dot{\psi} + \phi \cdot \dot{q} dt,$$

where $z = (p, q)$ and $\eta = (\phi, \psi)$ are elements in E . By analyzing in term of the Fourier series we easily obtain that

$$L(p, q) = (\bar{q}, -\bar{p}). \quad (2.37)$$

It is well known that

$$\begin{aligned} \exp(vL) &= 1 + vL + \frac{1}{2!}v^2L^2 + \frac{1}{3!}v^3L^3 + \frac{1}{4!}v^4L^4 + \cdots, \\ \cosh(vL) &= 1 + \frac{1}{2!}v^2L^2 + \frac{1}{4!}v^4L^4 + \cdots, \\ \sinh(vL) &= vL + \frac{1}{3!}v^3L^3 + \frac{1}{5!}v^5L^5 + \cdots. \end{aligned}$$

Thus, from (2.37) and recalling that $\bar{\bar{p}} = -p$ for every $p \in L^\vee$ one obtains

$$\exp(vL)(p, 0) = \cosh(v)(p, 0) - \sinh(v)(0, \bar{p}) \quad (2.38)$$

and in a similar way

$$\exp(vL)(0, q) = \cosh(v)(0, q) - \sinh(v)(\bar{q}, 0). \quad (2.39)$$

Adding (2.38) and (2.39) we have

$$\exp(vL)(p, q) = (\cosh(v)p - \sinh(v)\bar{q}, \sinh(v)\bar{p} + \cosh(v)q). \quad (2.40)$$

We can give an explicit formula for \hat{B} . Given $z \in X$ we write $z = (p, q) + z_0$, where $(p, q) \in E_0$ and z_0 . First we obtain a formula for $\hat{B}(p, q)$. From (2.8), (2.9), (2.40) and noting that $\bar{p} = q$ when $(p, q) \in E_-$ one has

$$B_1^{-1} \exp(vL) B_2(p, q) = (a_3 p, a_4 q), \quad (2.41)$$

where

$$a_3 = \frac{1}{\rho^{\beta-1}} (\cosh(v)\sigma^{\beta-1} - \sinh(v)\sigma^{\alpha-1})$$

and

$$a_4 = \frac{1}{\rho^{\alpha-1}} (-\sinh(v)\sigma^{\beta-1} + \cosh(v)\sigma^{\alpha-1}).$$

Using the formula for the projection into E_- given in Lemma 2.5 and Lemma 2.3 from (2.41) we obtain

$$\begin{aligned} \hat{B}(p, q) &= P_X(a_3 p, a_4 q) = \frac{1}{2}(a_3 p, a_4 q) - \frac{1}{2}L(a_3 p, a_4 q) \\ &= \frac{1}{2}(a_3 p, a_4 q) + \frac{1}{2}(a_4 p, a_3 q) = \frac{1}{2}(a_3 + a_4)(p, q) := \frac{1}{2}a_5(p, q). \end{aligned}$$

An easy computation shows that

$$a_5 = \left(\frac{\sigma^{\beta-1}}{\rho^{\beta-1}} + \frac{\sigma^{\alpha-1}}{\rho^{\alpha-1}} \right) \cosh(v) - \left(\frac{\sigma^{\alpha-1}}{\rho^{\beta-1}} + \frac{\sigma^{\beta-1}}{\rho^{\alpha-1}} \right) \sinh(v).$$

If we assume $\sigma > 1$ and $\rho < 1$ it is easy to see that $a_5 > 0$. Then we have

$$\hat{B}((p, q) + z_0) = \frac{1}{2}a_5(p, q) + \left(\frac{1}{\rho^{\beta-1}} p_0, \frac{1}{\rho^{\alpha-1}} q_0 \right)$$

so that \hat{B} is invertible. Thus, (I3) is satisfied. \square

Proof of Theorem 1.1. Using Lemmas 2.6–2.9, the hypothesis of Theorem 2.2 is satisfied. Thus a critical point z_K of J_K with critical value $c_K \geq \delta$ characterized by

$$c_K = \inf_{h \in \Gamma} \sup_{z \in Q} I(h(z, 1)). \quad (2.42)$$

Next we show that as a consequence of hypothesis (H4) for K large enough $\|z_K\|_\infty < K$ so that z_K is a solution of (1.1). Recalling that the constants σ and M are independent of K , from (2.7) we obtain that Q is a bounded set. Thus, there is a constants $c_8 > 0$ satisfying

that $\|z\| \leq c_8$ for every $z \in Q$. Then, from (H1), the definition of H_K , and the boundedness of Q , taking $h = id \in \Gamma$ from (2.42) we obtain

$$c_K \leq \sup_{z \in Q} J_K(z) \leq \frac{1}{2} \langle Lz, z \rangle \leq \frac{1}{2} c_8 \|L\|.$$

Denoting $z_K = (p_K, q_K)$ we have

$$\begin{aligned} \frac{1}{2} c_8 \|L\| &\geq c_K = J_K(z_K) = J_K(z_K) - J'_K(z_K) \cdot p_K \\ &= - \int_0^T (H_K(z_K, t) - H_K p(z_K, t) \cdot p_K) dt. \end{aligned} \quad (2.43)$$

Similarly, one has

$$\frac{1}{2} c_8 \|L\| \geq - \int_0^T (H_K(z_K, t) - H_K q(z_K, t) \cdot q_K) dt. \quad (2.44)$$

Multiplying (2.43) by $1/\alpha$, (2.44) by $1/\beta$, and adding we obtain

$$\begin{aligned} \left(\frac{1}{\alpha} + \frac{1}{\beta}\right) c_8 \|L\| &\geq - \left(\left(\frac{1}{\alpha} + \frac{1}{\beta}\right)\right) \int_0^T H_K(z_K, t) dt \\ &\quad + \int_0^T \frac{1}{\alpha} H_{Kp}(z, t) \cdot p_K + \frac{1}{\beta} H_{Kq}(z, t) \cdot q_K dt. \end{aligned} \quad (2.45)$$

Then, using hypothesis (H2) that is also satisfied by H_K with the same constants

$$\frac{1}{2} \left(\frac{1}{\alpha} + \frac{1}{\beta}\right) \|L\| c_8 \geq \left(1 - \left(\frac{1}{\alpha} + \frac{1}{\beta}\right)\right) \int_0^T H_K(z_K, t) dt. \quad (2.46)$$

Using Lemma 2.1 from (2.46) we obtain that for a constant c_9 independent of K ,

$$\|p_K\|_\alpha^\alpha + \|q_K\|_\beta^\beta \leq c_9. \quad (2.47)$$

Since z_K satisfies (1.1) we have from hypothesis (H4) and (2.45) that

$$\begin{aligned} \|\dot{z}_K\|_1 &= \int_0^T |H_{Kz}(z_K, t)| dt \\ &\leq a \int_0^T \left(\frac{1}{\alpha} H_{Kp}(z_K, t) \cdot p_K + \frac{1}{\beta} H_{Kq}(z_K, t) \cdot q_K\right) dt + Tb \leq c_{10} \end{aligned} \quad (2.48)$$

with c_{10} independent of K . Then (2.47) and (2.48) give that $\|z_K\|_\infty$ is bounded independent of K , which finishes the proof. \square

Proof of Theorem 1.2. We can take $T = 2\pi$ and choose $k \in \mathcal{N}$. It is convenient to make the change of variables $\tau = k^{-1}t$. Thus $z(t)$ is a $2k\pi$ periodic solution of (1.1), $\zeta(\tau) = z(k\tau)$ satisfies

$$\frac{d\zeta}{d\tau} = k\mathcal{J}H_z(\zeta, k\tau), \quad (2.49)$$

by

$$\frac{d\zeta(\tau)}{d\tau} = k \cdot \frac{dz(k\tau)}{dk\tau} = k\mathcal{J}H_z(z(k\tau), k\tau) = k\mathcal{J}H_z(\zeta(\tau), k\tau).$$

We seek a 2π periodic solution of (2.49). Since $kH(z, k\tau)$ satisfies (H0)–(H4), Theorem 1.1 provides a critical point $\zeta_{k,K}(\tau) \in E$ of

$$J_{k,K}(\zeta) = A(\zeta) - k \int_0^{2\pi} H_K(\zeta, k\tau) d\tau,$$

where K depends on k , which for appropriately large K is a classical solution of (2.49).

Let $\zeta_k(\tau)$ is a classical solution of (2.49). Note that $\zeta_1(k\tau)$ also satisfies (2.49) by

$$\frac{d\zeta_1(k\tau)}{d\tau} = k \cdot \frac{d\zeta_1(k\tau)}{dk\tau} = k \cdot \mathcal{J}H_z(\zeta_1(k\tau), k\tau).$$

Taking

$$J_k(\zeta) = A(\zeta) - k \int_0^{2\pi} H(\zeta(\tau), k\tau) d\tau,$$

if $\zeta_1(k\tau) = \zeta_k(\tau)$, we have

$$\begin{aligned} J_k(\zeta_k) &= A(\zeta_k(\tau)) - k \int_0^{2\pi} H(\zeta_k(\tau), k\tau) d\tau \\ &= \int_0^{2\pi} p_k(\tau) \cdot \frac{dq_k(\tau)}{d\tau} d\tau - k \int_0^{2\pi} H(\zeta_k(\tau), k\tau) d\tau \\ &= k \int_0^{2\pi} \left(p_1(k\tau) \cdot \frac{dq_1(k\tau)}{dk\tau} - H(\zeta_1(k\tau), k\tau) \right) d\tau \\ &= \int_0^{2k\pi} \left(p_1(s) \cdot \frac{dq_1(s)}{ds} - H(\zeta_1(s), s) \right) ds = kJ_1(\zeta_1). \end{aligned}$$

Thus, one has

$$d_k \equiv J_k(\zeta_k) = kJ_1(\zeta_1) = kd_1.$$

Since $d_1 > 0$ by Theorem 2.2, it follows that $d_k \rightarrow \infty$ as $k \rightarrow \infty$. We shall show next that this is impossible since d_k is bounded from above independently of k . Recall from Theorem 2.2, (2.16) and (H1) that

$$\begin{aligned} d_k &\leq \sup_{z \in Q} J_{k,K}(z) \\ &= \sup_{0 \leq s \leq \sigma, 0 \leq \|(p,q)+z_0\| \leq M} \left(\sigma^{\alpha+\beta-2} \left(\frac{1}{2} \lambda s^2 - \|(p,q)\|^2 \right) - k \int_0^{2\pi} H_K(z, k\tau) d\tau \right) \\ &\leq \lambda (\sigma(k))^{\alpha+\beta}, \end{aligned}$$

where we have written $\sigma(k)$ to emphasize its dependence on k . The parameter $\sigma(k)$ was determined in (2.33) and (2.34) when $s = \sigma$. The corresponding equation satisfied by $\sigma(k)$ is

$$\frac{1}{2} \lambda \sigma^{\alpha+\beta} - k c_6 c^\alpha \sigma^{\alpha\beta} + 2k\pi c_2 \leq 0$$

or

$$\frac{1}{2} \lambda \sigma^{\alpha+\beta} - k c_6 c^\beta \sigma^{\alpha\beta} + 2k\pi c_2 \leq 0$$

for all $\sigma \geq \sigma(k)$. It follows that

$$\sigma(k) = \max \left\{ \left(\frac{\lambda}{k c_6 c^\alpha} \right)^{1/(\alpha\beta - (\alpha+\beta))}, \left(\frac{4\pi c_2}{c_6 c^\alpha} \right)^{1/(\alpha\beta)} \right\}$$

or

$$\sigma(k) = \max \left\{ \left(\frac{\lambda}{k c_6 c^\beta} \right)^{1/(\alpha\beta - (\alpha+\beta))}, \left(\frac{4\pi c_2}{c_6 c^\beta} \right)^{1/(\alpha\beta)} \right\},$$

which implies that $\{\sigma(k) \mid k \in N\}$ is bounded. Thus the critical values d_k are bounded and therefore there is k_1 such that $\zeta_1(k\tau) \neq \zeta_k(\tau)$ for all $k \geq k_1$. Reapplying what we have just shown to the 2π periodic function $k_1 H(z, k_1 \tau)$ it follows that there is a sequence of nonzero 2π periodic solutions $z_j(\tau)$ to

$$\frac{dz}{d\tau} = j k_1 \mathcal{J} H_z(z, j k_1 \tau) \quad (2.50)$$

with $z_j(\tau) \neq z_1(j\tau)$ for all $j \geq k_2$. Moreover, from the form of (2.50) and the corresponding variational problem, $z_j(\tau) = \zeta_{jk_1}(\tau)$ and $z_j(\tau) \neq \zeta_1(jk_1 \tau)$ for all $j \geq k_2$. It follows that we have a sequence

$$\zeta_1(t), \zeta_{k_1}\left(\frac{t}{k_1}\right), \zeta_{k_1 k_2}\left(\frac{t}{k_1 k_2}\right), \dots,$$

of distinct nonzero solutions of (1.1) and the proof is complete. \square

References

- [1] P.L. Felmer, Periodic solutions of “superquadratic” Hamiltonian systems, *J. Differential Equations* 102 (1993) 188–207.
- [2] Y.M. Long, X.J. Xu, Periodic solutions for a class of nonautonomous Hamiltonian systems, *Nonlinear Anal.* 41 (2000) 455–463.
- [3] S.L. Li, A.M. Szulkin, Periodic solutions for a class of nonautonomous Hamiltonian systems, *J. Differential Equations* 112 (1994) 226–238.
- [4] Y.M. Long, Periodic solutions of superquadratic Hamiltonian systems with bounded forcing terms, *Math. Z.* 203 (1990) 453–467.
- [5] P.H. Rabinowitz, On subharmonic solutions of Hamiltonian systems, *Comm. Pure Appl. Math.* 33 (1980) 609–633.
- [6] P.H. Rabinowitz, Periodic solutions of large norm of Hamiltonian systems, *J. Differential Equations* 50 (1983) 33–48.
- [7] P.H. Rabinowitz, Periodic solutions of Hamiltonian systems, *Comm. Pure Appl. Math.* 31 (1978) 157–184.
- [8] S.J. Li, M. Willem, Applications of local linking to critical point theory, *J. Math. Anal. Appl.* 189 (1995) 6–32.
- [9] V. Benci, P.H. Rabinowitz, Critical point theorems for indefinite functionals, *Invent. Math.* 52 (1979) 241–273.
- [10] C.G. Liu, Subharmonic solutions of Hamiltonian systems, *Nonlinear Anal.* 42 (2000) 185–198.
- [11] P. Rabinowitz, Multiple critical points of perturbed symmetric functionals, *Trans. Amer. Math. Soc.* 272 (1982) 753–769.
- [12] F. Antonacci, Existence of periodic solutions of Hamiltonian systems with potential indefinite in sign, *Nonlinear Anal.* 29 (1997) 1353–1364.
- [13] Y.M. Long, Multiple solutions of perturbed superquadratic second order Hamiltonian systems, *Trans. Amer. Math. Soc.* 311 (1989) 749–780.
- [14] M. Willem, Subharmonic oscillations of convex Hamiltonian systems, *Nonlinear Anal.* 9 (1985) 1303–1311.
- [15] I. Ekeland, H. Hofer, Subharmonics for convex nonautonomous Hamiltonian systems, *Comm. Pure Appl. Math.* 40 (1987) 1–36.
- [16] R. Michalek, G. Tarantello, Subharmonic solutions with prescribed minimal period for nonautonomous Hamiltonian systems, *J. Differential Equations* 72 (1988) 28–55.
- [17] M. Girardi, M. Matzeu, Dual Morse index estimates for periodic solutions of Hamiltonian systems in some nonconvex superquadratic cases, *Nonlinear Anal.* 17 (1991) 481–497.
- [18] J.Q. Liu, Z.Q. Wang, Remarks on subharmonics with minimal periods of Hamiltonian systems, *Nonlinear Anal.* 20 (1993) 803–821.
- [19] Y. Xu, Subharmonic solutions for convex nonautonomous Hamiltonian systems, *Nonlinear Anal.* 28 (1997) 1359–1371.
- [20] E.A. de Silva, Subharmonic solutions for subquadratic Hamiltonian systems, *J. Differential Equations* 115 (1995) 120–145.
- [21] J. Yang, Y. Zheng, Z. Li, Multiple subharmonic solutions of nonautonomous Hamiltonian system, *J. Math. Res. Exposition* 13 (1993) 215–221.
- [22] K. Tanaka, Homoclinic orbits in a first order superquadratic Hamiltonian system: convergence of subharmonic orbits, *J. Differential Equations* 94 (1991) 315–339.
- [23] C.G. Liu, The stability of subharmonic solutions for Hamiltonian systems, *J. Math. Anal. Appl.* 240 (1999) 491–504.
- [24] A. Fonda, M. Willem, Subharmonic oscillations of forced pendulum-type equations, *J. Differential Equations* 81 (1989) 215–220.
- [25] A. Fonda, A.C. Lazer, Subharmonic solutions of conservative systems with nonconvex potentials, *Proc. Amer. Math. Soc.* 115 (1992) 183–190.
- [26] A. Fonda, M. Ramos, Large-amplitude subharmonic oscillations for scalar second-order differential equations with asymmetric nonlinearities, *J. Differential Equations* 109 (1994) 354–372.
- [27] Q. Wang, Z.Q. Wang, J.Y. Shi, Subharmonic oscillations with prescribed minimal period for a class of Hamiltonian systems, *Nonlinear Anal.* 28 (1997) 1273–1282.

- [28] N.P. Cac, A.C. Lazer, On second order, periodic, symmetric, differential systems having subharmonics of all sufficiently large orders, *J. Differential Equations* 127 (1996) 426–438.
- [29] M.Y. Jiang, Subharmonic solutions of second order subquadratic Hamiltonian systems with potential changing sign, *J. Math. Anal. Appl.* 244 (2000) 291–303.
- [30] E. Serra, M. Tarallo, S. Terracini, Subharmonic solutions to second-order differential equations with periodic nonlinearities, *Nonlinear Anal.* 41 (2000) 649–667.
- [31] N. Hirano, Subharmonic solutions for second order differential systems, *J. Math. Anal. Appl.* 196 (1995) 266–281.
- [32] R.E. Edwards, *Fourier Series*, vol. 2, A Modern Introduction, second ed., Springer-Verlag, New York, 1982.